

I-HEDGE: Determining the optimum complementary sets of taxa for conservation using evolutionary isolation

Evelyn L. Jensen, Arne Ø. Mooers, Adalgisa Caccone, and Michael A. Russello

Table S1. Pairwise values of ϕ_{ST} among species, calculated using the Kimura 2-parameter model.

Each value is significant at $p < 0.001$, except between *darwini* and *becki*, which is significant at $p < 0.05$.

	<i>darwini</i>	<i>donfaustoi</i>	<i>hoodensis</i>	<i>porteri</i>	<i>becki</i>	<i>abingdoni</i>	<i>ephippium</i>	<i>chathamensis</i>	<i>vandenburghi</i>	<i>microphyes</i>	<i>nigra</i>	<i>vicina</i>
<i>darwini</i>	0.00											
<i>donfaustoi</i>	0.92	0.00										
<i>hoodensis</i>	0.91	1.00	0.00									
<i>porteri</i>	0.84	0.91	0.93	0.00								
<i>becki</i>	0.11	0.92	0.92	0.87	0.00							
<i>abingdoni</i>	0.92	0.98	0.98	0.91	0.93	0.00						
<i>ephippium</i>	0.90	0.96	0.96	0.84	0.91	0.95	0.00					
<i>chathamensis</i>	0.91	0.99	1.00	0.92	0.92	0.98	0.96	0.00				
<i>vandenburghi</i>	0.93	0.99	0.99	0.85	0.93	0.98	0.95	0.99	0.00			
<i>microphyes</i>	0.92	0.99	0.99	0.81	0.92	0.98	0.94	0.99	0.67	0.00		
<i>nigra</i>	0.86	0.94	0.96	0.70	0.88	0.94	0.89	0.95	0.90	0.88	0.00	
<i>vicina</i>	0.85	0.88	0.89	0.74	0.87	0.89	0.85	0.89	0.27	0.36	0.71	0.00