

Table S2. Summary of the ANOVA statistics in the fitted linear model for the different *A. phaeospermum* strains effect on fresh root:shoot ratio.

Control	Estimate	Std. error	<i>t</i> -value	<i>P</i> (> <i>t</i>)
(Intercept)	0.3392	0.2334	1.4530	0.1505
Strain 1	0.0177	0.3301	0.0540	0.9573
Strain 2	0.0585	0.3301	0.1770	0.8598
Strain 3	0.0618	0.3301	0.1870	0.8518
Strain 4	0.0453	0.3301	0.1370	0.8912
Strain 5	0.0739	0.3301	0.2240	0.8233
Strain 6	0.0083	0.4367	0.0190	0.9848
Strain 7	0.0570	0.3301	0.1730	0.8632
Strain 8	0.1924	0.4367	0.4410	0.6608
Drought	Estimate	Std. error	<i>t</i> -value	<i>P</i> (> <i>t</i>)
(Intercept)	1.6698	0.3301	5.0580	<.0001
Strain 1	-0.3612	0.4668	-0.7740	0.4415
Strain 2	-1.0452	0.4668	-2.2390	0.0282
Strain 3	-0.7106	0.4668	-1.5220	0.1323
Strain 4	-1.0125	0.4668	-2.1690	0.0334
Strain 5	-0.3775	0.4668	-0.8090	0.4214
Strain 6	-0.5702	0.5474	-1.0420	0.3011
Strain 7	-0.9423	0.4668	-2.0180	0.0472
Strain 8	-1.2077	0.5597	-2.1580	0.0343

Data under control and drought treatments are shown. Significant *P*-values are indicated in bold text. Std. error: standard error.