



**Figure. Dummy tendency graphs.** For normal condition, the logical parameters given in Table 2 have been shown through dummy tendency graphs. Along x-axes resource entities and y-axes generation/synthesis level of targeted entities are taken. Logically absence/presence of negative regulator/s as resource indicating it is actually active/inactive for downregulation of the targeted entities [1,2].



**Figure. Dummy tendency graphs.** For normal condition, the logical parameters given in Table 2 have been shown through dummy tendency graphs. Along x-axes resource entities and y-axes generation/synthesis level of targeted entities are taken. Logically absence/presence of negative regulator/s as resource indicating it is actually active/inactive for downregulation of the targeted entities [1,2].



**Figure. Dummy tendency graphs.** For normal condition, the logical parameters given in Table 2 have been shown through dummy tendency graphs. Along x-axes resource entities and y-axes generation/synthesis level of targeted entities are taken. Logically absence/presence of negative regulator/s as resource indicating it is actually active/inactive for downregulation of the targeted entities [1,2].

## References

- [1] Paracha RZ, Ahmad J, Ali A, Hussain R, Niazi U, Tareen SHK, et al. Formal modelling of toll like receptor 4 and jak/stat signalling pathways: insight into the roles of socs-1, interferon- $\beta$  and proinflammatory cytokines in sepsis. *PloS one*. 2014;9(9):e108466.
- [2] Saeed MT, Ahmad J, Baumbach J, Pauling J, Shafi A, Paracha RZ, et al. Parameter estimation of qualitative biological regulatory networks on high performance computing hardware. *BMC systems biology*. 2018;12(1):1–15.