Figure 3B The original blot of GAPDH in K562 cells




Figure 3B The original blot of Bcl-XL in K562 cells




Figure 3B The original blot of Bad in K562 cells




Figure 3B The original blot of Survivin in K562 cells




Figure 3B The original blot of GAPDH in MEG-01 cells




Figure 3B The original blot of Bcl-XL in MEG-01 cells




Figure 3B The original blot of Bad in MEG-01 cells




Figure 3B The original blot of Survivin in MEG-01 cells




Figure 4B The original blot of GAPDH in K562 cells




Figure 4B The original blot of hTERT in K562 cells




Figure 4B The original blot of c-MYC in K562 cells




Figure 4B The original blot of GAPDH in MEG-01 cells




Figure 4B The original blot of hTERT in MEG-01 cells




Figure 4B The original blot of c-MYC in MEG-01 cells




Figure 5A The original blot of GAPDH in K562 cells




Figure 5A The original blot of PI3K in K562 cells




Figure 5A The original blot of p-PI3K in K562 cells




Figure 5A The original blot of GAPDH in K562 cells




Figure 5A The original blot of AKT in K562 cells




Figure 5A The original blot of P-AKT in K562 cells




Figure 5A The original blot of mTOR in K562 cells

mTOR	250kDa	
	9	
	_	

Figure 5A The original blot of P-mTOR in K562 cells




Figure 5A The original blot of ERK in K562 cells




Figure 5A The original blot of P-ERK in K562 cells




Figure 5A The original blot of MAPK in K562 cells




Figure 5A The original blot of P-MAPK in K562 cells




Figure 5B The original blot of GAPDH in MEG-01 cells




Figure 5B The original blot of PI3K in MEG-01 cells




Figure 5B The original blot of p-PI3K in MEG-01 cells




Figure 5B The original blot of GAPDH in MEG-01 cells




Figure 5B The original blot of AKT in MEG-01 cells




Figure 5B The original blot of P-AKT in MEG-01 cells




Figure 5B The original blot of mTOR in MEG-01 cells




Figure 5B The original blot of P-mTOR in MEG-01 cells




Figure 5B The original blot of ERK in MEG-01 cells




Figure 5B The original blot of p-ERK in MEG-01 cells




Figure 5B The original blot of MAPK in MEG-01 cells




Figure 5B The original blot of P-MAPK in MEG-01 cells




Figure 6C The original blot of GAPDH in K562 cells




Figure 6C The original blot of Bcl-XL in K562 cells




Figure 6C The original blot of Bad in K562 cells




Figure 6C The original blot of Survivin in K562 cells




Figure 6C The original blot of GAPDH in MEG-01 cells



Figure 6C The original blot of Bcl-XL in MEG-01 cells




Figure 6C The original blot of Bad in MEG-01 cells




Figure 6C The original blot of Survivin in MEG-01 cells



Figure 7B The original blot of GAPDH in K562 cells




Figure 7B The original blot of hTERT in K562 cells




Figure 7B The original blot of C-MYC in K562 cells




Figure 7B The original blot of GAPDH in MEG-01 cells




Figure 7B The original blot of hTERT in MEG-01 cells




Figure 7B The original blot of c-MYC in MEG-01 cells




Figure 8A The original blot of GAPDH in K562 cells




Figure 8A The original blot of PI3K in K562 cells




Figure 8A The original blot of p-PI3K in K562 cells




Figure 8A The original blot of GAPDH in K562 cells




Figure 8A The original blot of AKT in K562 cells




Figure 8A The original blot of P-AKT in K562 cells




Figure 8A The original blot of mTOR in K562 cells




Figure 8A The original blot of P-mTOR in K562 cells




Figure 8A The original blot of ERK in K562 cells




Figure 8A The original blot of p-ERK in K562 cells




Figure 8A The original blot of MAPK in K562 cells




Figure 8A The original blot of P-MAPK in K562 cells




Figure 8B The original blot of GAPDH in MEG-01 cells




Figure 8B The original blot of PI3K in MEG-01 cells




Figure 8B The original blot of p-PI3K in MEG-01 cells




Figure 8B The original blot of GAPDH in MEG-01 cells




Figure 8B The original blot of AKT in MEG-01 cells




Figure 8B The original blot of P-AKT in MEG-01 cells




Figure 8B The original blot of mTOR in MEG-01 cells




Figure 8B The original blot of P-mTOR in MEG-01 cells




Figure 8B The original blot of ERK in MEG-01 cells




Figure 8B The original blot of P-ERK in MEG-01 cells




Figure 8B The original blot of MAPK in MEG-01 cells

