

Original image of Figure 1, panel (A).


Line 1: molecular protein marker

Line 2: homogenate of non-transformed E. coli BL21.

Line 3: homogenate of E. coli BL21 transformed with empty vector.

Line 4: homogenate of E. coli BL21 transformed with pCold-ApoLpIII.


The gel was Coomassie blue-stained



Original image of Figure 1, panel (B).

Line 1: molecular protein marker


Line 2: rApoLp-III purified and Coomassie blue-stained.



Original image of Figure 1, panel (C).

Line 1: molecular protein marker

Line 2: rApoLp-III purified and silver-stained.




Original image of Figure 2, panel (A).

Line 1: molecular protein marker

Line 2: : rApoLp-III purified.

Line 3: *G. mellonella* moth's homogenate

The gel was Coomassie blue-stained




**Original image of Figure 2,
panel (B)**

Line 1: molecular Protein
marker

Line 2: rApoLp-III purified

Line 3: *G. mellonella* moth's
homogenate

The membrane was incubate




**Original image of Figure 2,
panel (C)**

Line 1: molecular Protein
marker

Line 2: rApoLp-III purified

Line 3: *G. mellonella* moth's
homogenate

The membrane was incubated
with pre-immune serum and



Original image of Figure 2, panel (D)

Line 1: molecular Protein marker

Line 2: rApoLp-III purified

Line 3: *G. mellonella* moth's
homogenate

The membrane was incubated only
with second antibody