

Table S1. Secondary outcomes of the participants

	NC (n=660)	EOPE (n=19)	LOPE (n=57)	LOPE vs. EOPE	EOPE vs. NC	LOPE vs. NC	P-value
Maternal outcomes							
Postpartum hemorrhage or blood transfusion	30 (4.5)	3 (15.8)	7 (12.3)	0.704	0.060	0.021	0.007
Admission to intensive care unit	10 (1.5)	4 (21.1)	4 (7.0)	0.102	<0.001	0.020	<0.001
Chorioamnionitis or postpartum infection	68 (10.3)	5 (26.3)	11 (19.3)	0.528	0.044	0.037	0.014
Neonatal outcomes							
Neonatal asphyxia	35 (2.7)	7 (18.4)	1 (0.9)	<0.001	<0.001	0.356	<0.001
Ventilator support	112 (8.5)	13 (34.2)	2 (1.8)	<0.001	<0.001	0.011	<0.001
Neonatal hypoglycemia	144 (10.9)	9 (23.7)	19 (16.7)	0.334	0.031	0.063	0.013
Sepsis	75 (5.7)	7 (18.4)	6 (5.3)	0.019	0.006	0.853	0.014
Blood transfusion	50 (3.8)	7 (18.4)	2 (1.8)	0.001	0.001	0.429	0.001
Hemorrhage	14 (1.1)	1 (2.6)	0 (0.0)	0.250	0.348	0.619	0.297

Data are presented as n (%). Abbreviations: NC, Normal control; EOPE, early-onset preeclampsia; LOPE, late-onset preeclampsia.